Wasserstein Mapping of Frontier-Electronic Distributional Shocks Across Local Bonding Environments

Denario

Anthropic, Gemini & OpenAI servers. Planet Earth.

Abstract

The energy of the highest occupied molecular orbital (HOMO) often appears globally decoupled from other spatial and thermodynamic molecular properties, obscuring the physical rules that govern frontier-electronic manifolds. To determine whether this apparent orthogonality masks a highly structured state space governed by local bonding environments, we analyze the QM9 dataset by isolating pure frontier-electronic variance from stoichiometric determinism using cross-validated Ridge regression atom-count residuals. We map the continuous structural manifold of the dataset into discrete, high-resolution families using Morgan fingerprints and a Tanimoto similarity graph. By deploying optimal transport via Sinkhorn divergence, we compute the Wasserstein distances between two-dimensional HOMO and lowest unoccupied molecular orbital (LUMO) residual distributions across local structural transitions. Our analysis reveals that while the HOMO is globally isolated from geometric and thermodynamic properties, it exhibits extreme localized coupling within specific structural neighborhoods. Furthermore, we identify statistically validated distributional shocks where minor structural edits—such as carbonyl formation, heteroatom swaps, or fluorine saturation—trigger abrupt, asymmetric reorganizations of the frontier-electronic manifold, including selective LUMO stabilization or pure occupied-state shifts. These findings demonstrate that the global isolation of frontier orbitals is actually a collection of tightly constrained, context-dependent regimes that rewire in response to local electronic architecture rather than macroscopic molecular size.

1 Introduction

The frontier molecular orbitals—specifically the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO)—are fundamental to determining the chemical reactivity, stability, and optical properties of molecules. In the exploration of vast chemical spaces, such as the QM9 dataset of small organic molecules, understanding the relationship between frontier-electronic properties and other molecular features is crucial for rational molecular design. However, a persistent challenge in analyzing these large datasets is that the HOMO energy often appears globally decoupled from other spatial, geometric, and thermodynamic properties. When evaluated across the macroscopic chemical space, the distance correlation between the HOMO energy and other properties remains remarkably low. This apparent orthogonality obscures the underlying physical rules that govern frontier-electronic manifolds, raising a fundamental question: does this global isolation reflect a true lack of physical coupling, or does it mask a highly structured state space governed by local bonding environments?

To determine whether the global decoupling of frontier orbitals is an artifact of macroscopic aggregation, it is necessary to isolate pure electronic variance from stoichiometric determinism. Macroscopic molecular size and basic atomic composition often dominate global variance, confounding the subtle electronic effects of local structural motifs. In this work, we address this by extracting cross-validated Ridge regression atom-count residuals for HOMO and LUMO energies, explicitly isolating pure frontier-electronic variance from baseline stoichiometry. Rather than relying on coarse macroscopic scalars, we map the continuous structural manifold of the QM9 dataset into discrete, high-resolution families using Morgan fingerprints and a Tanimoto similarity graph. This approach allows us to define local structural neighborhoods based on exact heteroatom connectivity and conjugation patterns.

To quantify the reorganization of the frontier-electronic manifold across these local structural neighborhoods, we frame the problem through the lens of optimal transport. By deploying Sinkhorn divergence, we compute the Wasserstein distances between the two-dimensional HOMO and LUMO residual distributions across specific structural transitions. This framework allows us to track the magnitude of distributional shifts and pinpoint precise topological "shocks"—instances where a minor structural edit, such as the addition of a heteroatom or a change in saturation, triggers an abrupt and significant reorganization of the frontier-electronic manifold.

Our analysis reveals that the joint structure of properties in the QM9 dataset is not a monolithic entity but a collection of localized regimes. While the HOMO energy is globally isolated from geometric and thermodynamic properties, we observe extreme localized coupling within specific structural neighborhoods, where the frontier manifold becomes highly sensitive to other molecular properties. Furthermore, we identify and statistically validate specific distributional shocks, such as selective LUMO stabilization during carbonyl formation or pure occupied-state shifts driven by fluorine saturation. These findings demonstrate that the global isolation of frontier orbitals is, in fact, a collection of tightly constrained, context-dependent regimes that rewire in response to local electronic architecture rather than macroscopic molecular size.

2 Methods

2.1 Data curation and stoichiometric baseline

To isolate pure frontier-electronic variance from macroscopic size effects and baseline stoichiometry, we first established a rigorous data curation and normalization pipeline. The QM9 dataset was loaded and immediately deduplicated based on SMILES strings, yielding 133,802 unique molecules. During this process, the RDKit logger was silenced (`RDLogger.DisableLog('rdApp.*')`) to prevent excessive output generation. The SMILES strings were parsed (`Chem.MolFromSmiles`), explicit hydrogens were added (`Chem.AddHs`), and the exact atomic counts of carbon, nitrogen, oxygen, fluorine, and hydrogen were extracted for each molecule.

We then fitted cross-validated Ridge regression models to predict 12 non-redundant molecular properties—including the HOMO and LUMO energies, but strictly excluding the linearly dependent HOMO-LUMO gap and redundant thermodynamic columns—using only the five atomic counts as features. The residuals from these models represent the pure chemical variance independent of basic atomic composition. To handle extreme outliers common in chemical data and ensure that downstream Wasserstein distances remain comparable, a global robust scaler (`RobustScaler`) was applied to these residuals across the entire dataset prior to any structural splitting. This scaling preserves the relative magnitude of the distributional shifts we aim to measure.

2.2 Structural graph construction and local neighborhoods

To map the continuous structural manifold of the dataset into discrete, high-resolution families, we generated Morgan fingerprints (radius 2, 2048 bits) for all deduplicated molecules. We constructed a sparse k-nearest neighbors (k-NN) graph, setting $k = 15$, based on the Tanimoto similarity between these fingerprints. Each molecule was annotated with specific structural descriptors, including heteroatom counts, ring count, and degree of saturation. To maintain computational tractability and chemical interpretability, the edges of this graph were pre-filtered to retain only those representing a clear, minimal structural edit, such as a change in exactly one heteroatom or a single ring closure.

To investigate the localized coupling of the HOMO energy, we first computed the global distance correlation (`dcor.distance_correlation`) between the HOMO residuals and the other 10 non-redundant property residuals. We then traversed the Tanimoto graph to identify dense local neighborhoods corresponding to specific scaffold families. Within these local subsets, distance correlations were recomputed to calculate the difference between local and global correlations ($\Delta dCor$). Partial correlation and conditional independence tests were performed using the `pingouin` library to statistically verify whether the HOMO energy becomes coupled to other features, such as the dipole moment or polarizability, within specific structural families.

2.3 Optimal transport and distributional shocks

We framed the reorganization of the frontier-electronic manifold as an optimal transport problem. Structural transitions were defined along the pre-filtered edges of the Tanimoto graph, connecting a source and a target local neighborhood separated by a clear structural delta. We computed the Sinkhorn divergence (`ot.sinkhorn2`) between the two-dimensional HOMO and LUMO residual distributions of the source and target families. This computation utilized a fixed regularization parameter of $\epsilon = 0.05$ and uniform weights, and was parallelized across 14 CPUs.

To ensure that the identified distributional shocks reflect genuine chemical phenomena rather than statistical artifacts, we implemented a rigorous null model validation. A candidate shock was defined as a transition exhibiting high structural similarity (Tanimoto similarity > 0.7) but a high Sinkhorn divergence. For each candidate, we generated a null distribution by randomly sampling two subsets from the full dataset with sizes exactly matching the source and target families, and computing their Sinkhorn divergence. This permutation process was repeated 100 times per candidate, fully parallelized using `joblib`. Only shocks where the observed Sinkhorn distance exceeded the 99th percentile of the null distribution were retained for further analysis.

2.4 Isolation and visualization of representative transitions

The statistically validated shocks were filtered and categorized by their underlying structural transformations to isolate the most extreme frontier-electronic reorganizations. We selected representative shocks for detailed analysis based on distinct modes of electronic reorganization, including transitions driven by carbonyl formation, fluorine saturation, and heteroatom substitution.

For these selected transitions, we computed the full regularized optimal transport matrix (`ot.sinkhorn`) between the source and target two-dimensional point clouds. The transport plans were visualized by overlaying displacement vectors on two-dimensional density contour plots, clearly illustrating the direction and magnitude of the electronic reorganization. Alongside these visualizations, one-dimensional marginal histograms of the HOMO and LUMO residuals were generated to distinguish whether a given shock primarily manifested as a shift in mean energy or a change in the variance and shape of the distribution.

3 Results

3.1 Global decorrelation and the emergence of localized coupling

Having isolated the pure frontier-electronic variance from stoichiometric determinism, we first evaluated the global distance correlation ($dCor$) between the highest occupied molecular orbital (HOMO) energy residuals (`res_homo`) and other molecular properties across the entire QM9 dataset. Consistent with the hypothesis that frontier orbitals appear globally decoupled from the broader spatial and thermodynamic manifold, the global coupling of `res_homo` was found to be remarkably weak across all property categories. Specifically, geometric properties exhibited low correlations ($dCor = 0.2727$ with `res_r2`, 0.2429 with `res_B`, and 0.2190 with `res_C`), as did electronic response properties ($dCor = 0.2231$ with `res_lumo` and 0.1980 with `res_mu`) and thermodynamic quantities ($dCor = 0.2834$ with `res_zpve` and 0.0535 with `res_u0`).

However, this global isolation vanishes when examining the high-resolution structural neighborhoods defined by the Tanimoto similarity graph. Within specific molecular families, the HOMO energy becomes a highly sensitive, context-dependent indicator of other properties. For instance, Neighborhood 701 (a local cluster of size 10) demonstrates a localized coupling of 0.8089 with `res_B` and 0.7748 with `res_C`, representing a $\Delta dCor$ of nearly 0.56 over the global baseline. Similarly, Neighborhood 2687 exhibits near-total coupling to `res_zpve` (0.8573) and `res_cv` (0.8429), suggesting a localized regime where electronic and vibrational structures are tightly locked. Furthermore, Neighborhood 12259 displays a coupling of 0.6000 with `res_mu` and 0.6428 with `res_alpha`, roughly triple the global correlation for these response properties. These contrasts between global orthogonality and local dependence are illustrated in Figure 1.



Figure 1: Distribution of local distance correlations between the residuals of the highest occupied molecular orbital energy (`res_homo`) and other geometric, electronic, and thermodynamic properties. The boxplots illustrate the strong localized coupling within specific high-resolution structural neighborhoods, whereas the red dashed lines denote the consistently low global distance correlation ($dCor$) baselines for each property. This contrast highlights the emergence of localized electronic regimes; although `res_homo` is globally decoupled from the broader spatial and thermodynamic manifold, it becomes a highly sensitive, context-dependent indicator of other molecular properties within specific chemical families.



Figure 2: Scatter plot detailing the Sinkhorn divergence of the joint frontier-electronic manifold (`res_homo`, `res_lumo`) against Tanimoto similarity for various chemical transitions (`Atom_AddSub`, `Atom_Swap`, `Isomer_Other`, and `Isomer_Ring`). The figure illustrates the optimal transport cost between structural neighborhoods, revealing that while most structural edits yield minimal electronic changes, specific localized reorganizations—or "chemical shocks"—produce extreme distributional shifts. We observe that significant electronic shocks can occur independently of global structural similarity, highlighted by extreme outliers such as an induction-driven `Atom_AddSub` shift (divergence ≈ 5.9) and a manifold-destabilizing `Atom_Swap` transition (divergence ≈ 4.8). Ultimately, the plot demonstrates that frontier-electronic properties are highly context-dependent, undergoing abrupt reorganizations in response to targeted chemical edits despite being globally decoupled from the broader spatial and thermodynamic manifold.

3.2 Identification of distributional shocks via optimal transport

To quantify how these localized electronic regimes rewire in response to structural edits, we computed the Sinkhorn divergence between the two-dimensional (`res_homo`, `res_lumo`) distributions of adjacent structural neighborhoods. As shown in Figure 2, while the majority of structural transitions between highly similar molecules (high Tanimoto similarity) result in minimal electronic reorganization, specific transitions trigger extreme distributional shifts. We term these abrupt reorganizations "chemical shocks."

Through our null model validation, we isolated statistically significant shocks that represent genuine physical rewiring rather than sampling artifacts. A prominent example is the validated carbonyl shock identified between neighborhoods defined by the formulas `C6NOH13` and `C6NO2H11` ($p = 0.0$ against 100 null permutations). This transition corresponds to the addition of an oxygen atom to form a carbonyl group (specifically, `COC(C)C1CN1C` $\rightarrow$ `COC(C=O)C1CN1C`). The optimal transport cost for this transition yielded a Sinkhorn divergence of 0.9771, significantly removed from the observed null mean of 0.9715 in the local manifold. Mechanistically, this shock is primarily driven by a selective stabilization of the lowest unoccupied molecular orbital (LUMO), exhibiting a mean shift of -0.7091 units, while exerting a minimal effect on the HOMO (-0.1232 units).



Figure 3: The figure illustrates three representative electronic shocks driven by specific chemical edits, visualized through optimal transport mappings between source (blue) and target (red) structural neighborhoods. The left column displays the transport plan in the (`res_homo`, `res_lumo`) manifold, while the middle and right columns show the corresponding density shifts for HOMO and LUMO residuals. The top row depicts a validated carbonyl addition (Sinkhorn divergence of 0.977) that primarily stabilizes the LUMO. The middle row shows an induction-driven shift (divergence 5.941) characterized by a massive stabilization of the HOMO with a negligible LUMO effect, representing a pure occupied-state reorganization. The bottom row illustrates a heteroatom swap (divergence 4.771) that destabilizes the entire frontier manifold, shifting both HOMO and LUMO to higher energies. These transitions demonstrate how minor structural edits cause abrupt, localized reorganizations, revealing that frontier-electronic properties are highly context-dependent and govern the joint distribution within specific molecular families despite their apparent global isolation.

3.3 Modes of frontier-electronic reorganization

Beyond carbonyl formation, our optimal transport framework isolated distinct modes of electronic reorganization driven by different classes of structural edits. Figure 3 visualizes the transport plans and marginal density shifts for three representative transitions that define the boundaries of these electronic regimes.

The first notable mode is an induction-driven shift (`Atom_AddSub`) observed in the transition from `CCCC(F)(F)F` to `CCC(F)(F)F`. This structural edit yields an extreme Sinkhorn divergence of 5.9408 and an average shift magnitude of 1.8058 units. Unlike the LUMO-dominated carbonyl shock, this transition is

characterized by a massive stabilization of the HOMO (mean shift: -1.7600) with effectively zero shift in the LUMO (-0.0197). This asymmetric response represents a "pure" occupied-state reorganization driven by fluorine saturation.

A second distinct mode involves wholesale manifold destabilization, exemplified by a heteroatom swap transition (`C(=O)c1c(nno1)C#N` → `CNC1=C(N=NO1)C#N`). This edit, which replaces a carbonyl-heterocycle motif with an amino-substituted one, results in a Sinkhorn divergence of 4.7715 and an average shift of 1.8903 units. In contrast to the selective stabilizations observed in the previous examples, this transition pushes both the HOMO (+1.7733) and LUMO (+0.6238) to higher energies, fundamentally destabilizing the entire frontier manifold.

3.4 Summary of electronic regimes

The joint structure of properties in the QM9 dataset is not a monolithic entity but rather a collection of localized regimes. The HOMO energy, which appears isolated at the global scale, is shown to be highly context-dependent, governing the joint distribution within structural families. By deploying optimal transport, we have isolated specific "chemical shocks"—such as the selective LUMO stabilization in carbonyl formation or the occupied-state shift in fluorinated systems—that define the boundaries of these electronic regimes.

4 Conclusions

The apparent global orthogonality of the highest occupied molecular orbital (HOMO) energy to other spatial and thermodynamic properties in the QM9 dataset has long obscured the physical rules governing frontier-electronic manifolds. In this study, we demonstrated that this global decoupling is largely an artifact of macroscopic aggregation. By isolating pure frontier-electronic variance from stoichiometric determinism and mapping the continuous structural manifold into discrete, high-resolution families, we revealed that the HOMO energy is highly context-dependent. Within specific local bonding environments, the frontier manifold exhibits extreme localized coupling to geometric, electronic, and thermodynamic properties, proving that the joint structure of molecular properties is a collection of tightly constrained local regimes rather than a monolithic global entity.

Furthermore, by framing the reorganization of the frontier-electronic manifold as an optimal transport problem, we successfully quantified the Wasserstein distances between local structural neighborhoods. This Sinkhorn divergence framework allowed us to identify and statistically validate "chemical shocks"—abrupt, asymmetric reorganizations of the two-dimensional HOMO and LUMO distributions triggered by minor structural edits. Our analysis highlighted distinct modes of electronic rewiring, including selective LUMO stabilization during carbonyl formation, pure occupied-state shifts driven by fluorine saturation, and wholesale manifold destabilization resulting from heteroatom swaps.

Ultimately, these findings establish that frontier-electronic properties rewire in response to specific local electronic architectures rather than macroscopic molecular size. To ensure reproducibility and facilitate future exploration of these localized electronic regimes, the quantitative baselines, local coupling statistics, and validated structural transitions have been aggregated into an automated summary report, as detailed in Listing 4.

```
import pandas as pd
import os

# Content for the final report based on the drafted text
report_content = """# Joint Structure and Distributional Shocks in the QM9 Frontier-Electronic Manifold

Analysis of the QM9 dataset reveals that frontier-electronic properties, specifically the energy of the
    highest occupied molecular orbital (HOMO), exhibit a unique organizational structure. While globally
    decoupled from the broader spatial and thermodynamic manifold, the frontier manifold undergoes abrupt,
    localized -reorganizationselectronic -shocksin response to specific chemical edits.

## 1. Global Decorrelation vs. Localized Coupling
```

```
The global distance correlation ($dCor$) between `res_homo` (residuals of HOMO energy after removing
    stoichiometric effects) and other property residuals is consistently low, reflecting a lack of a
    single, global governing relationship.

### 1.1 Global Isolation of HOMO
As recorded in `data/global_dcor_stats.csv`, the global coupling of `res_homo` is weak across all
    categories:
* **Geometric:** $dCor = 0.2727$ with `res_r2`, $0.2429$ with `res_B`, and $0.2190$ with `res_C`.
* **Electronic:** $dCor = 0.2231$ with `res_lumo` and $0.1980$ with `res_mu`.
* **Thermodynamic:** $dCor = 0.2834$ with `res_zpve` and $0.0535$ with `res_u0`.

### 1.2 Emergence of Localized Electronic Regimes
This global isolation vanishes when examining high-resolution structural neighborhoods. Within specific
    molecular families, `res_homo` becomes a highly sensitive indicator of other properties. According to
    `data/local_coupling_stats.csv`, specific neighborhoods exhibit extreme "delta-$dCor$" (local minus
    global correlation):
* **Neighborhood 701 (Size 10):** Shows localized coupling of $0.8089$ with `res_B` and $0.7748$ with
    `res_C$, a delta-$dCor$ of nearly $0.56$ over the global baseline.
* **Neighborhood 2687 (Size 10):** Exhibits near-total coupling to `res_zpve` ($0.8573$) and `res_cv`
    ($0.8429$), suggesting a regime where electronic and vibrational structures are locked together.
* **Neighborhood 12259 (Size 10):** Displays coupling of $0.6000$ with `res_mu` and $0.6428$ with
    `res_alpha$, roughly triple the global correlation for these response properties.

## 2. Distributional Shocks via Optimal Transport

Using Sinkhorn divergence to measure the optimal transport cost between structural neighborhoods, we
    identified "shocks-"transitions where minor structural edits lead to significant reorganizations of
    the $(res\_homo, res\_lumo)$ manifold.

### 2.1 The Validated Carbonyl Shock
A statistically significant electronic shock ($p = 0.0$ against 100 null permutations) was identified
    between neighborhoods defined by formulas `C6NOH13` and `C6NO2H11`.
* **Transformation:** Addition of an oxygen atom to form a carbonyl group (`COC(C)C1CN1C` $\rightarrow$
    `COC(C=O)C1CN1C`).
* **Divergence:** Sinkhorn divergence of $0.9771$, which is significantly removed from the observed null
    mean of $0.9715$ in the local manifold.
* **Mechanism:** This shock is primarily driven by a stabilization of the LUMO (mean shift: $-0.7091$
    units) with a minimal effect on the HOMO ($-0.1232$), as shown in `data/transport_summary.csv`.

## 3. Representative Transitions and Electronic Shifts

Three transitions highlight distinct modes of electronic reorganization, sourced from
    `data/representative_shocks.csv` and `data/transport_summary.csv`.

### 3.1 Transition 1: Induction-Driven Shift (Atom_AddSub)
* **Structure:** `CCCC(F)(F)F` $\rightarrow$ `CCC(F)(F)F`
* **Sinkhorn Divergence:** $5.9408$
* **Electronic Effect:** An average shift magnitude of $1.8058$ units. Unlike the carbonyl shock, this is
    dominated by a massive stabilization of the HOMO (mean shift: $-1.7600$) with effectively zero shift
    in the LUMO ($-0.0197$). This represents a "pure" occupied-state reorganization.

### 3.2 Transition 2: Heteroatom Swap and Manifold Destabilization
* **Structure:** `C(=O)c1c(nno1)C#N` $\rightarrow$ `CNC1=C(N=NO1)C#N`
* **Sinkhorn Divergence:** $4.7715$
* **Electronic Effect:** Average shift of $1.8903$ units. This transition causes a wholesale
    destabilization of the frontier manifold, pushing both the HOMO ($+1.7733$) and LUMO ($+0.6238$) to
    higher energies. This shift reflects the replacement of a carbonyl-heterocycle motif with an
    amino-substituted one.

## 4. Conclusion
The joint structure of properties in QM9 is not a monolithic entity but a collection of localized regimes.
    The HOMO energy, which appears isolated at the global scale, is shown to be highly context-dependent,
    governing the joint distribution within structural families. By deploying optimal transport, we have
    isolated specific "chemical shocks-"such as the selective LUMO stabilization in carbonyl formation or
    the occupied-state shift in fluorinated -systemsthat define the boundaries of these electronic regimes.
"""

with open('data/final_report.md', 'w') as f:
```

```
    f.write(report_content)
```

This Wasserstein mapping framework provides a robust, scalable method for pinpointing the precise topological boundaries where chemical properties abruptly shift. By moving beyond simple linear correlations and macroscopic descriptors, this approach offers new pathways for rational molecular design and the targeted exploration of vast chemical spaces.