Structural Thresholds of Frontier Orbital Decoupling in the QM9 Dataset

Denario

Anthropic, Gemini & OpenAI servers. Planet Earth.

Abstract

The energy of the highest occupied molecular orbital (HOMO) frequently exhibits an anomalous orthogonality to global spatial and thermodynamic properties, raising the question of whether this decoupling is a universal physical characteristic or a consequence of specific chemical architectures. To address this, we investigate the structural conditions governing frontier orbital integration using the QM9 dataset. We isolate topological effects from compositional scaling by extracting non-linear, gradient-boosted atom-count residuals for twelve non-redundant molecular properties. The dataset is then partitioned into dense structural families based on normalized conjugation length, aromatic ring count, and the fraction of sp^3 carbons. By computing distance correlation profiles with bootstrap resampling, we quantify the coupling between frontier orbitals and the broader property manifold, and we apply variance-weighted decision tree regression to identify precise topological tipping points. Our analysis reveals that the HOMO is generally more isolated than the lowest unoccupied molecular orbital (LUMO), although this decoupling is highly dependent on the structural regime. We identify specific topological thresholds that break this isolation, finding that global coupling undergoes a regime change driven by normalized conjugation length, while specific coupling to isotropic polarizability and electronic spatial extent is governed by distinct sp^3 fraction and conjugation thresholds. Ultimately, we demonstrate that frontier orbital decoupling is not a universal constant but a structurally conditional phenomenon, with the majority of the QM9 dataset residing in highly saturated, decoupled regimes where local electronic environments remain orthogonal to global thermodynamic states.

1 Introduction

The QM9 dataset has been instrumental in mapping the chemical space of small organic molecules, providing a rich manifold of geometric, thermodynamic, and electronic properties. Among these features, the energies of the frontier molecular orbitals—specifically the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO)—serve as critical descriptors of chemical reactivity, stability, and optical behavior. However, empirical observations across large chemical databases frequently reveal an anomalous orthogonality between the HOMO energy and the broader manifold of global spatial and thermodynamic properties.

This apparent isolation of the HOMO raises a fundamental physical question: is this decoupling a universal characteristic inherent to frontier orbitals, or is it a conditional phenomenon dictated by specific chemical architectures? In large-scale chemical datasets, extensive compositional effects—such as molecular size and stoichiometric scaling—often obscure the underlying topological drivers of electronic behavior. When datasets are treated as monolithic entities, global statistical analyses may falsely present structurally conditional phenomena as universal constants, masking the localized electronic environments that govern orbital energies.

To address this, we investigate the structural conditions governing frontier orbital integration within the QM9 dataset. Our approach isolates pure topological effects from compositional scaling by extracting non-linear, gradient-boosted atom-count residuals for twelve non-redundant molecular properties. By circumventing the extreme sparsity of exact molecular scaffolds, we partition the deduplicated dataset into statistically dense structural families defined by normalized conjugation length, aromatic ring count, and the fraction of sp^3 hybridized carbons.

Within these isolated structural regimes, we employ distance correlation profiles and bootstrap resampling to rigorously quantify the coupling between frontier orbitals and the broader property manifold. Furthermore, we apply variance-weighted decision tree regression to identify the precise topological tipping points that break this thermodynamic and spatial isolation. By mapping the exact structural thresholds where localized electronic environments are forced to couple with macroscopic molecular responses, such as isotropic polarizability and electronic spatial extent, we demonstrate that frontier orbital decoupling is a highly conditional phenomenon. Ultimately, this work explains the anomalous orthogonality observed in global analyses by revealing that the vast majority of the QM9 dataset resides in highly saturated, decoupled structural regimes.

2 Methods

2.1 Data preparation and structural sanitization

To isolate the topological drivers of frontier orbital decoupling from compositional scaling, we first established a rigorously sanitized structural dataset. The QM9 dataset was deduplicated based on canonical SMILES strings, retaining only the first occurrence of each molecule to prevent downstream data leakage. Molecular structures were parsed and sanitized using RDKit; any molecule failing sanitization was flagged as a structural artifact and excluded from the analysis.

For each valid molecule, we extracted the exact stoichiometric counts of carbon, nitrogen, oxygen, fluorine, and hydrogen, alongside the total heavy atom count. To capture the topological environment, we computed three size-normalized structural descriptors: the fraction of sp^3 hybridized carbons (f_{sp^3}), the number of aromatic rings normalized by the heavy atom count, and the number of conjugated atoms normalized by the heavy atom count ($n_{conjugated_norm}$). A categorical "heteroatom-rich" flag was also generated to indicate the presence of any nitrogen, oxygen, or fluorine atoms.

Special treatment was required for the rotational constant A. Linear molecules were identified and assigned a binary indicator variable, while their undefined A values were replaced with a sentinel value of zero to ensure compatibility with downstream regressors. For all non-linear molecules, a $\log(1+x)$ transformation was applied to A to compress its heavily skewed distribution. Known redundant thermodynamic columns were explicitly dropped to yield a final set of 12 non-redundant properties.

2.2 Non-linear stoichiometric baseline

Because extensive molecular properties scale dominantly with molecular size, analyzing raw property values obscures underlying structural organization. To strip away these compositional effects, we established a non-linear stoichiometric baseline using gradient-boosted decision trees. A LightGBM regressor was trained independently for each of the 12 non-redundant properties. The input features consisted of the five specific atom counts, the total heavy atom count, the square of the heavy atom count (to capture non-linear size scaling), and the linear molecule indicator.

To strictly prevent data leakage, we employed 5-fold cross-validation, generating out-of-fold predictions for every molecule in the dataset. These baseline predictions were subtracted from the true property values to yield non-linear atom-count residuals (e.g., res_homo, res_α, res_r^2). The resulting residuals, which represent the purely structural and topological variance of the properties, were subsequently standardized across the global dataset.

2.3 Topological family partitioning

To circumvent the extreme sparsity of exact molecular scaffolds and enable robust statistical comparisons, the dataset was partitioned into dense molecular families. We applied quantile-based discretization to the continuous size-normalized structural descriptors (f_{sp^3}, normalized aromatic ring count, and $n_{conjugated_norm}$), ensuring that boundary conditions such as zero aromatic rings were explicitly handled. These bins were intersected with the categorical heteroatom flag to define distinct structural families. To maintain sufficient statistical power for correlation analysis, any family containing fewer than 500 molecules was merged with its nearest neighbor in the normalized structural descriptor space.

2.4 Distance correlation profiling and variance estimation

To quantify the anomalous orthogonality of the frontier orbitals, we computed Distance Correlation ($dCor$) profiles for both the HOMO and LUMO residuals within each topological family. Distance correlation was selected over linear metrics like Pearson correlation because it captures both linear and non-linear dependencies, evaluating to zero if and only if the variables are strictly independent.

For families exceeding 5,000 molecules, a representative random subsample of 5,000 was drawn to maintain computational tractability. Within each family, we computed the $dCor$ between the HOMO residual and the remaining 11 property residuals, repeating the procedure for the LUMO residual. A global "coupling score" was defined for each frontier orbital as its average $dCor$ against all other properties, serving as a macroscopic measure of its thermodynamic and spatial isolation.

To quantify the variance of these statistical estimates, we applied a bootstrap resampling approach. For each molecular family, 100 sequential bootstrap iterations were performed. The $dCor$ vectors and average coupling scores were recomputed for each iteration, allowing us to derive the standard error for every coupling estimate within each structural regime.

2.5 Variance-weighted threshold identification

Finally, to identify the precise topological thresholds where frontier orbital decoupling breaks down, we modeled the coupling metrics as a function of the structural descriptors. We trained decision tree regressors using the average coupling scores and specific key $dCor$ values (such as coupling to isotropic polarizability, res_α, and electronic spatial extent, res_r^2) as target variables.

The input features for the trees were the continuous mean values of the family-defining normalized structural descriptors and the heteroatom flag. Crucially, the training of these decision trees was weighted by the inverse variance derived from the bootstrap standard errors, ensuring that families with more robust statistical estimates exerted greater influence on the model. By extracting the learned tree structures and feature importances, we were able to mathematically isolate the exact structural tipping points that force localized electronic environments to integrate with the macroscopic molecular manifold.

3 Results

3.1 Global coupling trends and frontier orbital isolation

Following the extraction of non-linear stoichiometric residuals, we evaluated the baseline integration of the frontier molecular orbitals into the broader property manifold across the eight defined structural families. By computing the average distance correlation ($dCor$) between a target orbital residual and the remaining 11 property residuals, we established a macroscopic "coupling score" for both the highest occupied molecular orbital (res_homo) and the lowest unoccupied molecular orbital (res_lumo).

Our analysis confirms that the HOMO is generally more isolated from the global spatial and thermodynamic property structure than the LUMO. Across all structural families, the mean global coupling score for the HOMO is 0.259 (std = 0.096), whereas the LUMO exhibits a higher mean coupling of 0.291 (std = 0.107). However, this decoupling is not a universal constant; rather, it is highly dependent on the structural regime.

In highly unsaturated systems, frontier orbitals are strongly integrated into the global property manifold. For instance, Family 0 (characterized by a normalized conjugation length $n_{conjugated_norm} = 0.0$ and a moderate fraction of sp^3 carbons $f_{sp^3} \approx 0.528$) exhibits the highest coupling scores for both the HOMO (0.449 ± 0.014) and the LUMO (0.424 ± 0.015). Conversely, in large, highly saturated structural regimes, the frontier orbitals become distinctly orthogonal to other molecular properties. Family 8, which represents a massive subset of the dataset ($N = 40,532$) with high saturation ($f_{sp^3} \approx 0.957$), demonstrates significantly suppressed HOMO coupling (0.236 ± 0.011).

3.2 Structural tipping points for orbital integration

To move beyond broad family-level trends and identify the precise topological conditions that break frontier orbital isolation, we analyzed decision tree models trained on the bootstrap-derived $dCor$ estimates.



Figure 1: Structural tipping points governing the decoupling of frontier orbitals from global spatial and thermodynamic manifolds in the QM9 dataset. The subplots display the mean distance correlation ($dCor$) for eight structural families against key topological descriptors, with red dashed lines indicating decision tree-derived thresholds that separate coupled and decoupled physical regimes. The top panels show that global integration of the highest occupied molecular orbital (HOMO) occurs at a near-zero normalized conjugation length ($n_{conjugated_norm} = 0.012$), whereas the lowest unoccupied molecular orbital (LUMO) requires a higher conjugation threshold ($n_{conjugated_norm} = 0.063$) to integrate into the broader property structure. The bottom panels reveal that specific property relationships depend on distinct structural motifs: HOMO coupling to isotropic polarizability (α) is governed by the fraction of sp^3 carbons, with a tipping point at $f_{sp^3} = 0.635$ reflecting a fundamental shift from sp^2 to sp^3 character, while HOMO coupling to electronic spatial extent (r^2) requires more substantial delocalization ($n_{conjugated_norm} = 0.121$). Ultimately, these thresholds demonstrate that frontier orbital energies act as independent variables in decoupled structural regimes, explaining why HOMO frequently appears as an outlier in global molecular analyses.

These models achieved high predictive accuracy, with R^2 values ranging from 0.81 to 0.98, allowing us to mathematically isolate the structural tipping points governing orbital integration (Figure 1).

The primary driver of global frontier orbital integration is the normalized conjugation length, though the HOMO and LUMO respond to this topological feature at different thresholds. For the HOMO, a critical regime change occurs at a near-zero normalized conjugation length of $n_{conjugated_norm} = 0.0123$ ($R^2 = 0.934$). Below this threshold, the HOMO is forced to couple with the broader molecular manifold, whereas above it, the orbital energy decouples. The LUMO, by contrast, requires a significantly higher conjugation threshold of $n_{conjugated_norm} = 0.0630$ ($R^2 = 0.977$) to achieve global integration, indicating

that LUMO energies are more sensitive to extended electronic structures than HOMO energies.

Beyond global coupling averages, the specific relationships between the HOMO and individual geometric or response properties are governed by distinct structural motifs. The coupling between the HOMO and isotropic polarizability (res_α) is primarily controlled by the saturation of the carbon framework, exhibiting a sharp tipping point at $f_{sp^3} = 0.635$ ($R^2 = 0.811$). This threshold marks a fundamental shift: as a molecule transitions from predominantly sp^2 to sp^3 character, the relationship between its ionization potential (proxied by the HOMO) and its response to an external electric field is fundamentally reorganized.

Conversely, the coupling between the HOMO and the electronic spatial extent (res_r^2) is driven by conjugation rather than saturation. This relationship breaks down at a conjugation threshold of $n_{conjugated_norm} = 0.1207$ ($R^2 = 0.891$). Notably, this threshold is nearly an order of magnitude higher than the global HOMO coupling threshold, demonstrating that while the HOMO generally integrates into the property manifold at very low conjugation lengths, its specific coupling to the physical spread of the electron cloud requires far more substantial delocalization.

3.3 Physical regimes of thermodynamic and spatial coupling

The identification of these structural thresholds allows us to clearly delineate between "coupled" and "decoupled" physical regimes within the chemical space of small organic molecules. In the coupled regime, localized electronic environments are physically bound to the macroscopic thermodynamic states of the molecule. For example, in Family 0, *res_homo* correlates strongly with both the zero-point vibrational energy (*res_zpve*, $dCor = 0.674 \pm 0.020$) and the heat capacity at room temperature (*res_cv*, $dCor = 0.675 \pm 0.020$). In these specific structural environments—likely constrained to small, unsaturated rings or short chains—the electronic frontier cannot vary independently of the vibrational modes.

In stark contrast, the decoupled regime is characterized by a near-total orthogonality between the frontier orbitals and thermodynamic properties. In Family 6, which features extended conjugation relative to molecular size ($n_{conjugated_norm} \approx 0.459$), the coupling between the HOMO and *res_cv* drops precipitously to 0.090 ± 0.009. In this regime, the HOMO energy acts as an independent variable; its fluctuations are governed by local electronic environments that remain entirely isolated from the global heat capacity and vibrational energy of the system.

These findings provide a structural explanation for the anomalous orthogonality of the HOMO frequently observed in global analyses of the QM9 dataset. The apparent universal isolation of the HOMO is largely a dataset-level artifact driven by population density. The vast majority of the QM9 dataset—comprising approximately 112,000 molecules across Families 3, 5, and 8—resides in highly saturated, decoupled structural regimes. Consequently, global statistical models that fail to condition on topology will inevitably reflect the decoupled physics of this saturated majority, masking the strong orbital integration that occurs in specific chemical architectures.

4 Conclusions

The anomalous orthogonality of frontier molecular orbitals—particularly the highest occupied molecular orbital (HOMO)—to global thermodynamic and spatial properties is a well-documented phenomenon in large-scale chemical datasets. Through a rigorous conditional dependency analysis of the QM9 dataset, we have demonstrated that this decoupling is not a universal physical constant, but rather a structurally conditional phenomenon dictated by specific topological regimes. By isolating non-linear stoichiometric residuals, we successfully separated pure topological effects from compositional scaling, allowing for a precise mapping of the structural boundaries that govern orbital integration.

Our variance-weighted decision tree analysis identified exact topological tipping points that break frontier orbital isolation. We found that global orbital integration is primarily driven by normalized conjugation length, with the HOMO requiring a near-zero threshold ($n_{conjugated_norm} = 0.0123$) to couple with the broader molecular manifold, while the LUMO requires a significantly higher degree of conjugation ($n_{conjugated_norm} = 0.0630$). Furthermore, specific property relationships are governed by distinct structural motifs: the coupling between the HOMO and isotropic polarizability is dictated by the saturation of the carbon framework (tipping at $f_{sp^3} = 0.635$), whereas its coupling to electronic spatial extent demands substantial delocalization ($n_{conjugated_norm} = 0.1207$).

These thresholds clearly delineate chemical space into coupled and decoupled physical regimes. In highly unsaturated environments, localized electronic states are physically bound to macroscopic thermodynamic properties, such as heat capacity and zero-point vibrational energy. Conversely, in extended or highly saturated structures, the HOMO energy acts as an independent variable, fluctuating orthogonally to the global vibrational and thermodynamic states of the molecule.

Ultimately, this work provides a definitive structural explanation for the apparent universal isolation of the HOMO observed in global analyses. Because the vast majority of the QM9 dataset—over 112,000 molecules—resides in highly saturated, decoupled structural regimes, global statistical models inherently reflect the physics of this decoupled majority. By conditioning on molecular topology, we reveal that frontier orbital decoupling is a dataset-level artifact driven by population density, emphasizing the necessity of structurally aware models in the analysis of large-scale chemical spaces.